

## anti- HDAC10 antibody

### Product Information

Catalog No.:	FNab10065
Size:	100µg
Form:	liquid
Purification:	Immunogen affinity purified
Purity:	≥95% as determined by SDS-PAGE
Host:	Rabbit
Clonality:	polyclonal
Clone ID:	None
IsoType:	IgG
Storage:	PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12 months (Avoid repeated freeze / thaw cycles.)

### Background

Responsible for the deacetylation of lysine residues on the N-terminal part of the core histones(H2A, H2B, H3 and H4). Histone deacetylation gives a tag for epigenetic repression and plays an important role in transcriptional regulation, cell cycle progression and developmental events. Histone deacetylases act via the formation of large multiprotein complexes.

### Immunogen information

Immunogen:	histone deacetylase 10
Synonyms:	HD10, HDAC10, histone deacetylase 10
Observed MW:	68 kDa
Uniprot ID :	Q969S8

### Application

Reactivity:	Human
Tested Application:	ELISA, IHC
Recommended dilution:IHC:	1:20-1:200

### Wuhan Fine Biotech Co., Ltd.


B9 Bld, High-Tech Medical Devices Park, No. 818 Gaoxin Ave. East Lake High-Tech Development Zone. Wuhan, Hubei, China(430206)

Tel : (0086)027-87384275

Fax: (0086)027-87800889

[www.fn-test.com](http://www.fn-test.com)

Image:



Immunohistochemistry of paraffin-embedded human colon cancer tissue slide using FNab10065(HDAC10 Antibody) at dilution of 1:200